

Spin magnetic susceptibility in the two-layer Hubbard model

Ferdinando Mancini ^{a,b}, Volodymyr Turkowski ^{a,b,1}

^a *Dipartimento di Scienze Fisiche "E.R. Caianiello" e Unit  I.N.F.M. di Salerno
Universit  di Salerno, 84081 Baronissi (SA), Italy*

^b *IIASS "E.R. Caianiello", 84019 Vietri sul Mare (SA), Italy*

Abstract

The two-layer Hubbard model is studied by using the composite operator method. Magnetic properties of the model in the normal state are analyzed by studying the uniform static spin magnetic susceptibility as a function of doping and temperature. The relevance of the model to describe the physical behavior of two-layered cuprate high- T_c superconductors is discussed.

Keywords: Magnetic susceptibility; Two-plane Hubbard model; YBCO

1. Introduction

The theoretical description of high- T_c superconductors (HTSC) still remains one of the most difficult and important tasks of the modern solid-state physics. There is a widespread opinion that anti-ferromagnetic correlations in the $Cu - O$ planes play an important role in the behavior of these systems. In this sense it is interesting to study the magnetic properties of HTSC in the framework of some strongly correlated electronic model. It was shown (see, for example, Refs.[1,2]), that the magnetic normal properties of the lanthanum family materials can be qualitatively described by using the 2D Hubbard model in the composite operator framework.

The aim of this paper is to study such properties, namely the uniform static spin magnetic suscepti-

bility, in the case of the two-plane Hubbard model which, we believe, can describe the behavior of the two-layered yttrium family materials.

2. Results

The two-plane Hubbard model is described by the next Hamiltonian

$$H = \sum_{i,j,\sigma} (t_{ij} - \mu \delta_{ij}) c_{\sigma}^{\dagger}(i) c_{\sigma}(j) + U \sum_i n_{\uparrow}(i) n_{\downarrow}(i).$$

Here we use the standard notations; the nearest hopping constant takes two values t and t_z for intra- and inter- plane hopping, correspondingly.

The composite operator method calculations give the next expression for the spin magnetic susceptibility in the static approximation

$$\chi(\mathbf{k}, \omega) = \frac{2}{n^2 - n - 2D} [n(Q_{1111}^R + 2Q_{1112}^R + Q_{1212}^R) + (2 - n)(Q_{1212}^R + 2Q_{1222}^R + Q_{2222}^R)],$$

¹ Corresponding author. E-mail: vturk@sa.infn.it




Fig. 1. The values of χ_0 are plotted as a function of n for different values of t_z at $U/t = 4$ and $T/t = 0.01$.

where n is the particle density and D is the double occupancy.

The function $Q_{\alpha\beta\gamma\delta}^R(\mathbf{k}, \omega)$ is the retarded part of the Fourier transformed expression of $Q_{\alpha\beta\gamma\delta}(i, j) = G_{\alpha\beta}(i, j)G_{\gamma\delta}(j, i)$. The thermal causal Green's functions (GFs) $G(i, j)$ are defined by $G(i, j) = \langle T[\Psi(i)\Psi^\dagger(j)] \rangle$ in the fermionic basis $\Psi^\dagger(i) = (\xi^\dagger(i), \eta^\dagger(i))$, where $\xi(i) = [1 - n(i)]c(i)$ and $\eta(i) = n(i)c(i)$ are the Hubbard's operators.

As it was shown previously (e.g. [3]), the single particle retarded GFs can be determined self-consistently with requirement to satisfy the Pauli principle $\langle \xi(i)\eta^\dagger(i) \rangle = 0$.

In this paper we present the calculations of the uniform static spin magnetic susceptibility $\chi_0 = \chi(\mathbf{k} \rightarrow 0, \omega = 0)$. In Fig.1 the values of χ_0 are plotted versus n for different values of the hopping constant t_z . Susceptibility χ_0 demonstrates a maximum at some value of n_c which depends on t_z . As in the 2D case, this peak is a consequence of the van Hove singularity (vHs) in the density of states. Numerical studies show that the value of n_c increases almost linearly with t_z and $n_c = 1$ at $t_z \simeq 1.4$ in the case $U/t = 4$ and $T/t = 0.01$. The experimental results [4] also demonstrate the maximum for $\chi_0(n)$ in $YBa_2Cu_3O_{7-x}$.

In Fig.2 χ_0 as a function of temperature is presented for different values of doping. Qualitatively the same behavior for χ_0 was found experimentally for some double-layered materials [4,5]. As it also follows from this picture, χ_0 demonstrate a sharp maximum at $T = 0$ when $n = n_c$.




Fig. 2. The values of χ_0 are plotted as a function of T for different values of n at $U/t = 4$ and $t_z = 1$.

It should be mentioned that the position of the vHs also changes with t' in the two-dimensional $t - t' - U$ Hubbard model [6,7]. However, the composite operator studies show [8] that for obtaining the right critical value n_c , in accordance with experiments on ittrium family materials, t' should be taken positive ~ 0.5 , whereas it is impossible with positive values to reproduce the right form of the Fermi surface after the ARPES data [9]. In this sense, from the composite operator method point of view, the two-layer Hubbard model looks more promising for describing the normal properties of the ittrium family materials.

References

- [1] F. Mancini, S. Marra, H. Matsumoto, Physica C 252 (1995) 361.
- [2] F. Mancini, H. Matsumoto and D. Villani, Phys. Rev. B 57 (1998) 6143.
- [3] A. Avella, F. Mancini, D. Villani *et al.*, Int. J. Mod. Phys. B 12 (1998) 81.
- [4] D.C. Johnson, S.K. Sinha, A.J. Jacobson and J.M. Newsam, Physica C 153-155 (1988) 572.
- [5] R.J. Cava, B. Batlogg, R.B. van Dover *et al.*, Phys. Rev. Lett. 58 (1987) 1676.
- [6] Q. Si, Yu. Zha, K. Levin, J.P. Lu, Phys. Rev. B 47 (1993) 9055.
- [7] P. Benard, L.Chen, and A.-M.S. Trembley, Phys. Rev. B 47 (1993) 15217.
- [8] A. Avella, F. Mancini, D. Villani, and H. Matsumoto, cond-mat/9707088.
- [9] R. Liu *et al.*, Phys. Rev. B 45 (1992) 5614.