

# Elementary evaluations of some Euler sums

Donal F. Connon

## Abstract

This short note contains elementary evaluations of some Euler sums.

It was shown by the author in [1, Eq. (4.4.155zi)] that

$$(-1)^{p+1} n \int_0^1 (1-t)^{n-1} \log^p t \, dt = p! \sum_{k=1}^n \binom{n}{k} \frac{(-1)^k}{k^p}$$

and with  $p = 1$  we obtain

$$n \int_0^1 (1-t)^{n-1} \log t \, dt = \sum_{k=1}^n \binom{n}{k} \frac{(-1)^k}{k} = -H_n^{(1)}$$

where  $H_n^{(r)}$  are the generalised harmonic numbers  $H_n^{(r)} = \sum_{k=1}^n \frac{1}{k^r}$ .

The summation then gives us

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^q} &= - \sum_{n=1}^{\infty} \frac{1}{n^{q-1}} \int_0^1 (1-t)^{n-1} \log t \, dt \\ &= - \int_0^1 \sum_{n=1}^{\infty} \frac{(1-t)^{n-1}}{n^{q-1}} \log t \, dt \end{aligned}$$

and we obtain

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^q} = - \int_0^1 \frac{Li_{q-1}(1-t) \log t}{1-t} dt$$

where  $Li_s(x)$  are the polylogarithm functions  $Li_s(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^s}$ .

With  $q = 2$  we have

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^2} = -\int_0^1 \frac{Li_1(1-t) \log t}{1-t} dt = \int_0^1 \frac{\log^2 t}{1-t} dt$$

With integration by parts we have

$$\int \frac{\log^2 t}{1-t} dt = 2Li_3(t) - 2Li_2(t) \log t - \log(1-t) \log^2 t + c$$

and we end up with the well-known result originally derived by Euler

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^2} = 2\zeta(3)$$

With  $q = 3$  we have

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^3} = -\int_0^1 \frac{Li_2(1-t) \log t}{1-t} dt$$

and we see that

$$\int \frac{Li_2(1-t) \log t}{1-t} dt = \frac{1}{2} [Li_2(1-t)]^2 + c$$

This then gives us the known result

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^3} = \frac{1}{2} \zeta^2(2)$$

Georgiou and Philippou [3] derived the following formula in 1983

$$\sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^{2p+1}} = \frac{1}{2} \sum_{j=2}^{2p} (-1)^j \zeta(j) \zeta(2p-j+2)$$

and we may therefore deduce the corresponding integral

$$\int_0^1 \frac{Li_{2p}(1-t) \log t}{1-t} dt = \frac{1}{2} \sum_{j=2}^{2p} (-1)^j \zeta(j) \zeta(2p-j+2)$$

We also have the product

$$\left[ H_n^{(1)} \right]^2 = n^2 \int_0^1 (1-t)^{n-1} \log t \, dt \int_0^1 (1-u)^{n-1} \log u \, du$$

and we get the summation

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\left[ H_n^{(1)} \right]^2}{n^q} &= \sum_{n=1}^{\infty} \frac{1}{n^{q-2}} \int_0^1 (1-t)^{n-1} \log t \, dt \int_0^1 (1-u)^{n-1} \log u \, du \\ &= \int_0^1 \int_0^1 \frac{Li_{q-2}[(1-t)(1-u)] \log t \log u \, du \, dt}{(1-t)(1-u)} \end{aligned}$$

Letting  $q = 2$  we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\left[ H_n^{(1)} \right]^2}{n^2} &= \int_0^1 \int_0^1 \frac{Li_0[(1-t)(1-u)] \log t \log u \, du \, dt}{(1-t)(1-u)} \\ &= \int_0^1 \int_0^1 \frac{\log t \log u \, du \, dt}{1 - (1-t)(1-u)} \end{aligned}$$

The Wolfram Integrator instantly gives us

$$\int \frac{\log t}{1 - (1-t)(1-u)} \, dt = \frac{1}{1-u} \log t \log \left( \frac{1 - (1-t)(1-u)}{u} \right) + \frac{1}{1-u} Li_2 \left( -\frac{(1-u)t}{u} \right) + c$$

and hence we have

$$\int_0^1 \frac{\log t}{1 - (1-t)(1-u)} \, dt = \frac{1}{1-u} Li_2 \left( -\frac{(1-u)}{u} \right)$$

We then obtain

$$\sum_{n=1}^{\infty} \frac{\left[ H_n^{(1)} \right]^2}{n^2} = \int_0^1 \frac{\log u}{1-u} Li_2 \left( -\frac{(1-u)}{u} \right) \, du$$

and it may be noted that the Wolfram Integrator cannot evaluate this integral.

We now employ the identity originally obtained by Landen [4] in 1780

$$Li_2\left(-\frac{(1-u)}{u}\right) = -\frac{1}{2}\log^2 u - Li_2(1-u)$$

with the result that

$$\int_0^1 \frac{\log u}{1-u} Li_2\left(-\frac{(1-u)}{u}\right) du = -\frac{1}{2} \int_0^1 \frac{\log^3 u}{1-u} du - \int_0^1 \frac{Li_2(1-u) \log u}{1-u} du$$

It is well known that  $\int_0^1 \frac{\log^3 u}{1-u} du = -6\zeta(4)$  and the second integral has already been evaluated above. We accordingly get

$$\int_0^1 \frac{\log u}{1-u} Li_2\left(-\frac{(1-u)}{u}\right) du = \frac{17}{4} \zeta(4)$$

and hence we have

$$\sum_{n=1}^{\infty} \frac{[H_n^{(1)}]^2}{n^2} = \frac{17}{4} \zeta(4)$$

as originally discovered by de Doelder [2] in 1991.

Letting  $q = 3$  we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{[H_n^{(1)}]^2}{n^3} &= \int_0^1 \int_0^1 \frac{Li_1[(1-t)(1-u)] \log t \log u}{(1-t)(1-u)} dt du \\ &= - \int_0^1 \int_0^1 \frac{\log[1-(1-t)(1-u)] \log t \log u}{(1-t)(1-u)} dt du \end{aligned}$$

The Wolfram Integrator can evaluate the integral

$$\int \frac{\log[1-(1-t)(1-u)] \log t}{(1-t)(1-u)} dt$$

but additional work is required to check if the definite integral exists.

## References:

[1] D.F. Connon, Some series and integrals involving the Riemann zeta function, binomial

coefficients and the harmonic numbers, Volume IV, 2007. [arXiv:0710.4028](https://arxiv.org/abs/0710.4028)

- [2] P.J. de Doelder, On some series containing  $\psi(x) - \psi(y)$  and  $(\psi(x) - \psi(y))^2$  for certain values of  $x$  and  $y$ . J. Comput. Appl. Math. 37, 125-141, 1991.
- [3] C. Georghiou and A.N. Philippou, Harmonic sums and the Zeta function. Fibonacci Quart., 21, 29-36, 1983.
- [4] J. Landen, Mathematical Memoirs, 1, 1780.